arXiv:hep-ex/0506071v1 28 Jun 2005

Charm and Beauty Production at HERA–B

Ulrich Husemann for the HERA–B Collaboration

Fachbereich Physik, Universität Siegen, D–57068 Siegen, Germany
now at University of Rochester, Rochester, New York 14627

Abstract. The HERA–B experiment at DESY has acquired a data-set of approximately 300,000 decays $\mathrm{J}/\psi \to \ell^+\ell^-$ during its 2002/2003 data-taking period. These data are used to analyze the production of heavy quarks in proton-nucleus interactions at a center-of-mass energy of 41.6 GeV. In this article, preliminary results of two measurements are discussed, a measurement of nuclear effects in the production of J/ψ mesons and a measurement of the $\mathrm{b}\overline{\mathrm{b}}$ production cross section.

HERA–B: DETECTOR, TRIGGER AND DATA-SET

The HERA–B detector, depicted in Fig. 1, is a fixed-target spectrometer with large angular acceptance. Protons from the halo of the HERA proton beam are collided with an internal wire target. The target consists of up to eight thin wires of different materials, distributed in two target stations. Each wire can be moved independently to adjust the interaction rate.

The HERA–B detector consists of the following sub-detectors: A silicon micro-strip vertex detector is used to reconstruct and separate primary and secondary vertices. The tracking detectors comprise micro-strip gaseous chambers with gas electron multiplier foils in the inner acceptance region of high particle flux and honeycomb drift chambers in the outer region. Particle identification is performed by a ring-imaging Čerenkov counter, an electromagnetic calorimeter, and a four-layer muon detector.

Lepton pairs from J/ψ decays are enriched by a multi-level trigger system. The first trigger level is implemented as a hardware track trigger in the outer tracking detector. Starting points for the track search are provided by pretriggers in the muon detector and the calorimeter. At the second trigger level, a software trigger running on a PC farm, track candidates are extrapolated to the vertex detector, and a two-prong vertex fit is performed. Accepted events are reconstructed online.

The HERA–B experiment has taken data between October 2002 and February 2003. The data sample recorded with the lepton pair trigger contains approximately 150,000 decays into each of the channels $\mathrm{J}/\psi \to \mathrm{e}^+\mathrm{e}^-$ and $\mathrm{J}/\psi \to \mu^+\mu^-$, 90,000 of which were taken with two target wires of different materials simultaneously. Based on this data set, the HERA–B collaboration has analyzed the production of charmonium states such as J/ψ, $\psi(2S)$ and χ_c, as well as the production of open and hidden beauty, e.g. $\mathrm{b}\overline{\mathrm{b}}$ and Υ.



FIGURE 1. Elevation view of the HERA–B detector.

NUCLEAR EFFECTS IN J/ψ PRODUCTION

Many theoretical models of J/ψ production in proton-nucleus interactions predict modifications of the production cross section by nuclear effects, e.g. due to the absorption of final state particles in the nuclear environment [1, 2], or due to coherent proton-nucleus scattering [3, 4]. The nuclear effects are commonly parametrized by the power law

$$\sigma_{\mathrm{p}A} = \sigma_{\mathrm{p}N} \cdot A^{\alpha(x_{\mathrm{F}},p_{\mathrm{T}})}. \tag{1}$$

Here, $\sigma_{\mathrm{p}A}$ is the proton-nucleus cross section, $\sigma_{\mathrm{p}N}$ is the proton-nucleon cross section, and A is the atomic mass number of the target. A value of the nuclear suppression parameter $\alpha(x_{\mathrm{F}}, p_{\mathrm{T}})$ smaller than unity corresponds to the suppression of J/ψ production. The cross section σ is related to the number of detected particles N, the integrated luminosity $\mathscr{L}$, and the detection efficiency ε by $\sigma = N/(\mathscr{L}\varepsilon)$. Using this relation for a tungsten target and a carbon target operated at the same time, Eq. (1) can be solved for α:

$$\alpha = \frac{1}{\log\left(A_{\mathrm{W}}/A_{\mathrm{C}}\right)} \cdot \log\left(\frac{N_{\mathrm{W}}}{N_{\mathrm{C}}} \cdot \frac{\mathscr{L}_{\mathrm{C}}}{\mathscr{L}_{\mathrm{W}}} \cdot \frac{\varepsilon_{\mathrm{C}}}{\varepsilon_{\mathrm{W}}}\right). \tag{2}$$

The measurement hence includes measuring three ratios, the ratio of J/ψ yields, the ratio of luminosities per wire, and the ratio of efficiencies. The analysis thus benefits from the advantages of a relative measurement, in which many systematic uncertainties cancel out. The J/ψ yields are determined by fits to the J/ψ invariant mass spectrum in bins of the kinematic variables. To measure the luminosities, the number of primary vertices per target wire is counted in data recorded with a zero-bias trigger in parallel to the lepton pair trigger. A detailed Monte Carlo (MC) simulation of the HERA–B detector and trigger is performed to calculate the ratio of efficiencies. Details of the analysis in the decay channel $\mathrm{J}/\psi \to \mu^+\mu^-$ can be found in [5].

The measured nuclear suppression parameter α in the decay channel $\mathrm{J}/\psi \to \mu^+\mu^-$ is shown in Fig. 2 as a function of x_{F} and p_{T}. The data show good agreement with previous measurements in the overlap region and extend the existing data to negative x_{F}. In the covered range of $-0.375 \leq x_{\mathrm{F}} < 0.125$, a small constant suppression of J/ψ production is observed, with an average suppression parameter of $\overline{\alpha} = 0.969 \pm 0.003(\mathrm{stat.}) \pm 0.021(\mathrm{syst.})$.



FIGURE 2. Nuclear suppression parameter α in the decay channel $J/\psi \rightarrow \mu^+\mu^-$ as a function of x_F (left) and p_T (right), compared to previous measurements by the E866 [6] and the NA50 [7] collaborations. All data points include statistical and systematic uncertainties.

THE B$\overline{\text{B}}$ PRODUCTION CROSS SECTION

At the HERA–B center-of-mass energy of 41.6 GeV, beauty production is close to its kinematic threshold. This results in large theoretical uncertainties due to the resummation of soft gluons and increased sensitivity to the mass of the b quark. Previous measurements of the $b\bar{b}$ production cross section in a similar energy range suffered from small statistics and large systematic uncertainties.

In HERA–B, the $b\bar{b}$ production cross section is measured in the inclusive decay channel $b\bar{b} \rightarrow J/\psi X \rightarrow \ell^+\ell^- X$, where $\ell = e, \mu$. The average flight distance of B mesons in the HERA–B detector amounts to approximately 8 mm. Therefore, B candidate events can be selected based on the separation of the J/ψ decay vertex from the primary vertex on the target wire. The best separation from background decays such as prompt J/ψ decays and semileptonic b and c decays is achieved by a combined cut on the significance of the J/ψ detached vertex and the impact parameter to the target wire. Furthermore, combinatorial background is controlled by comparing the background upstream and downstream of the target with respect to the proton flight direction. A fit to the B lifetime and the requirement of a third track at the J/ψ decay vertex serve as independent confirmations of the B flavor. For the final $b\bar{b}$ cross section result, the consistent results in the dielectron and the dimuon channels using the 2002/2003 datasample are combined with the previously published result [8] from the 2000 data-taking.

To minimize the influence of external input, the result of the measurement is first presented relative to the J/ψ production cross section and restricted to the kinematic acceptance of the HERA–B detector:

$$R_{b\bar{b}} = \frac{\Delta\sigma_{b\bar{b}}}{\Delta\sigma_{J/\psi}} = \varepsilon\, \frac{N_{b\bar{b}\rightarrow J/\psi X}}{N_{\text{prompt } J/\psi}} = 0.033 \pm 0.005(\text{stat.}) \pm 0.004(\text{syst.}). \qquad (3)$$

Here, $\Delta\sigma_{b\bar{b}}$ and $\Delta\sigma_{J/\psi}$ are the production cross sections for $b\bar{b}$ and J/ψ within the HERA–B acceptance, $N_{b\bar{b}\rightarrow J/\psi X}$ and $N_{\text{prompt } J/\psi}$ are the number of $b\bar{b}$ candidates and the number of prompt J/ψ. The efficiency parameter ε includes the branching fractions,



FIGURE 3. Preliminary cross section for $b\bar{b}$ production as a function of the proton energy, compared to theoretical predictions [12, 13] and a previous HERA–B measurement based on the 2000 data-set [8].

the efficiency of the detached vertex selection and the nuclear dependences for $b\bar{b}$ and J/ψ events. The efficiencies are determined from a detailed MC simulation. The result is extrapolated to the full phase-space with the help of PYTHIA [9]. Since to date, there is no published HERA–B measurement of the J/ψ production cross section, a value of $\sigma_{J/\psi} = 352 \pm 2(\text{stat.}) \pm 26(\text{syst.})\,\text{nb}/\text{nucl.}$ is assumed [8]. This value is obtained by averaging $\sigma_{J/\psi}$ as measured by the E789 and E771 experiments [10, 11] and extrapolating the result to HERA–B energies. Note that a significant upward revision of this value is anticipated from the HERA–B measurement. The resulting $b\bar{b}$ production cross section is depicted in Fig. 3, along with previous measurements and theoretical predictions. The preliminary numerical result is $\sigma_{b\bar{b}} = 9.9 \pm 1.5(\text{stat.}) \pm 1.4(\text{syst.})\,\text{nb}/\text{nucl.}$

ACKNOWLEDGMENTS

The author would like to thank the DIS05 crew for the well-organized and interesting conference. This work was supported by the German Bundesministerium für Bildung und Forschung under the contract number 5HB1PEA/7. Special thanks to the HERA–B charmonium and beauty working groups for their help in preparing this conference contribution and to DESY for the kind financial support.

REFERENCES

1. R. Vogt, *Phys. Rev.*, **C61**, 035203 (2000).
2. R. Vogt, *Nucl. Phys.*, **A700**, 539–554 (2002).
3. B. Z. Kopeliovich, A. Tarasov, and J. Hüfner, *Nucl. Phys.*, **A696**, 669–714 (2001).
4. K. G. Boreskov and A. B. Kaidalov, *JETP Lett.*, **77**, 599–602 (2003).
5. U. Husemann, *Measurement of Nuclear Effects in the Production of* J/ψ *Mesons with the HERA–B Detector*, Ph.D. thesis, Universität Siegen, Germany (2005), DESY-THESIS-2005-05.
6. M. J. Leitch et al., *Phys. Rev. Lett.*, **84**, 3256–3260 (2000).
7. B. Alessandro et al., *Eur. Phys. J.*, **C33**, 31–40 (2004).
8. I. Abt et al., *Eur. Phys. J.*, **C26**, 345–355 (2003).
9. T. Sjöstrand, *Comput. Phys. Commun.*, **82**, 74–90 (1994).
10. M. H. Schub et al., *Phys. Rev.*, **D52**, 1307–1315 (1995).
11. T. Alexopoulos et al., *Phys. Lett.*, **B374**, 271–276 (1996).
12. R. Bonciani, S. Catani, M. L. Mangano, and P. Nason, *Nucl. Phys.*, **B529**, 424–450 (1998).

13. N. Kidonakis and R. Vogt, *Eur. Phys. J.*, **C36**, 201–213 (2004).